

Etiopatogènia i tractament dels trastorns mentals greus

Publicacions 2010

Sumatori factor d'impacte : 104,663

Ahuir M, Bernardo M, Serna E, Ochoa S, Carlson J, Escartín G, Gutiérrez-Zotes A, Valero J, Salamero M, Cañizares S, Fernández Egea E, Cañete J, Gallo P. Adaptación y validación española de la Escala Pronóstica para la Esquizofrenia de Strauss y Carpenter.. *Rev Psiquiatr Salud Mental* 2010 ;2 (4) :150-9

Barajas A, Baños I, Ochoa S, Usall J, Huerta E, Dolz M, Sánchez B, Villalta Gil V, Foix A, Haro JM. Gender differences in incipient psychosis.. *Eur J Psychiatry* 2010 ;24 (3):176-194 Factor d'impacte: 0,462 Q4

Borges G, Nock MK, Haro JM, Hwang I, Sampson NA, Alonso J, Andrade LH, Angermeyer MC, Beautrais A, Bromet E, Bruffaerts R, De Girolamo G, Florescu S, Gureje O, Hu C, Karam EG, Kovess V, Lee S, Levinson D, Medina Mora ME, Ormel J, Posada Villa J, Sagar R, Tomov T, Uda H, Williams DR, Kessler RC. Twelve-month prevalence of and risk factors for suicide attempts in the world health organization world mental health surveys.. *J Clin Psychiatry* 2010 ;71 (12) :1617-28 Factor d'impacte: 5,218Q1

Brébion G, Laroí F, Van der Linden M. Associations of hallucination proneness with free-recall intrusions and response bias in a nonclinical sample.. *J Clin Exp Neuropsychol* 2010 ;32 (8) :847-54 Factor d'impacte: 1,885 Q3

Bruffaerts R, Demyttenaere K, Borges G, Haro JM, Chiu WT, Hwang I, Karam EG, Kessler RC, Sampson N, Alonso J, Andrade LH, Angermeyer MC, Benjet C, Bromet E, De Girolamo G, De Graaf R, Florescu S, Gureje O, Horiguchi I, Hu C, Kovess V, Levinson D, Posada Villa J, Sagar R, Scott K, Tsang A, Vassilev SM, Williams DR, Nock MK. Childhood adversities as risk factors for onset and persistence of suicidal behaviour.. *Br J Psychiatry* 2010 ;197 (1) :20-7 Factor d'impacte: 5,777 Q1

Casas E, Escandell MJ, Ribas M, Ochoa S. Instrumentos de evaluación en rehabilitación psicosocial.. *Rev As Esp Neuropsiquiatría* 2010 ;30 (105) :25-47

De Graaf R, Radovanovic M, Van Laar M, Fairman B, Degenhardt L, Aguilar Gaxiola S, Bruffaerts R, De Girolamo G, Fayyad J, Gureje O, Haro JM, Huang Y, Kostychenko S, Lépine JP, Matschinger H, Medina Mora ME, Neumark Y, Ormel J, Posada Villa J, Stein DJ, Tachimori H, Wells JE, Anthony JC. Early Cannabis Use and Estimated Risk of Later Onset of Depression Spells: Epidemiologic Evidence From the Population-based World Health Organization World Mental Health Survey Initiative.. *Am J Epidemiol* 2010 ;172 (2) :149-59 Factor d'impacte: 5,589 Q1

De Portugal E, González N, Miriam V, Haro JM, Usall J, Cervilla JA. Gender differences in delusional disorder: Evidence from an outpatient sample.. *Psychiatry Res* 2010;177 (1-2) :235-9 Factor d'impacte: 2,373 Q2

Escuder G, Gomollón LG, Ochoa S, Ramos MJ, González Cáceres A, Stuart T. Validación clínica de la escala Camberwell Assessment of Need-Forensic version, CANFOR para la evaluación de necesidades de personas con trastorno mental en el ámbito penitenciario.. *Actas Esp Psiquiatr* 2010 ;38 (3) :129-137 Factor d'impacte: 0,547Q4

Fuentealba P, Klausberger T, Karayannis T, Suen WY, Huck J, Tomioka R, Rockland K, Capogna M, Studer M, Morales M, Somogyi P. Expression of COUP-TFII nuclear receptor in

restricted GABAergic neuronal populations in the adult rat hippocampus.. *J Neurosci* 2010 ;30 (5) :1595-609 Factor d'impacte: 7,178 Q1

Fullana MA, Vilagut G, Rojas S, Mataix D, De Graaf R, Demyttenaere K, Haro JM, De Girolamo G, L epine JP, Matschinger H, Alonso J. ESEMed/MHEDEA 2000 investigators. Obsessive-compulsive symptom dimensions in the general population: Results from an epidemiological study in six European countries.. *J Affect Disord* 2010 ;124 (3) :291-9 Factor d'impacte: 3,763 Q1

Gabilondo A, Rojas S, Vilagut G, Haro JM, Fern andez A, Pinto Meza A, Alonso J. Epidemiology of major depressive episode in a southern European country: Results from the ESEMed-Spain project.. *J Affect Disord* 2010 ;120 (1-3) :76-85 Factor d'impacte: 3,763 Q1

Gonz alez Pinto A, Reed C, Novick D, Bertsch J, Haro JM. Assessment of Medication Adherence in a Cohort of Patients with Bipolar Disorder.. *Pharmacopsychiatry* 2010 ;43(7) :263-70 Factor d'impacte: 2,317 Q2

Hong J, Reed C, Novick D, Haro JM, Windmeijer F, Knapp M. The Cost of Relapse for Patients with a Manic/Mixed Episode of Bipolar Disorder in the EMBLEM Study.. *Pharmacoeconomics* 2010 ;28 (7) :555-66 Factor d'impacte: 2,612 Q2

Kessler RC, Birnbaum HG, Shahly V, Bromet E, Hwang I, McLaughlin KA, Sampson N, Andrade LH, De Girolamo G, Demyttenaere K, Haro JM, Karam AN, Kostyuchenko S, Kovess V, Lara C, Levinson D, Matschinger H, Nakane Y, Browne MO, Ormel J, Posada Villa J, Sagar R, Stein DJ. Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO World Mental Health Survey Initiative.. *Depress Anxiety* 2010 ;27 (4) :351-64 Factor d'impacte: 2,926 Q2

Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, Aguilar Gaxiola S, Alhamzawi AO, Alonso J, Angermeyer MC, Benjet C, Bromet E, Chatterji S, De Girolamo G, Demyttenaere K, Fayyad J, Florescu S, Gal G, Gureje O, Haro JM, Hu C, Karam EG, Kawakami N, Lee S, L epine JP, Ormel J, Posada Villa J, Sagar R, Tsang A, Ust un TB, Vassilev SM, Viana MC, Williams DR. Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys.. *Br J Psychiatry* 2010 ;197 (5) :378-85 Factor d'impacte: 5,777 Q1

Kovess V, Dezetter A, De Graaf R, Haro JM, Bruffaerts R, Briffault X, Gilbert F, Codony M, Alonso J. ESEMed/MHEDEA 2000 investigators. Religious advisors' role in mental health care in the European Study of the Epidemiology of Mental Disorders survey.. *Soc Psychiatry Psychiatry Epidemiol* 2010 ;45 (10) :989-98 Factor d'impacte: 2,052 Q3

Levinson D, Lakoma MD, Petukhova M, Schoenbaum M, Zaslavsky AM, Angermeyer MC, Borges G, Bruffaerts R, De Girolamo G, De Graaf R, Gureje O, Haro JM, Hu C, Karam AN, Kawakami N, Lee S, L epine JP, Browne MO, Okoliyski M, Posada Villa J, Sagar R, Viana MC, Williams DR, Kessler RC. Associations of serious mental illness with earnings: results from the WHO World Mental Health surveys.. *Br J Psychiatry* 2010 ;197 :114-21 Factor d'impacte: 5,777 Q1

Monta n es F, Gastaminza X, Catal a M, Ruiz Sanz F, Ruiz L azaro P, Herreros O, Garc a Giral M, Ortiz Guerra J, Alda J, Mojarro D, Cant o T, Mardomingo M, Sasot J, P amias M, Rey F. GEITDAH. Consenso del GEITDAH sobre el trastorno por d eficit de atenci n/hiperactividad.. *Rev Neurol* 2010 ;51 (10) :633-37 Factor d'impacte: 1,234 Q3

Moreno C, Merch an J,  lvarez M, Baeza I, Alda J, Mart inez Cantarero C, Parellada M, S anchez B, De la Serna E, Gir aldez M, Arango C. Metabolic effects of second-generation antipsychotics in bipolar youth: comparison with other psychotic and nonpsychotic diagnoses.. *Bipolar Disord* 2010 ;12 (2) :172-84 Factor d'impacte: 5,502 Q1 D1

Navarra J, Alsius A, Soto S, Spence C. Assessing the role of attention in the audiovisual integration of speech.. *Inform Fusion* 2010 ;11 (1) :4-11 Factor d'impacte: 2,212 Q1

- Navarra J, Alsius A, Velasco I, Toso Faraco S, Spence C. Perception of audiovisual speech synchrony for native and non-native language.. *Brain Res* 2010 ;1323 :84-93 Factor d'impacte: 2,463 Q3
- Novick D, Reed C, Haro JM, González Pinto A, Perrin E, Aguado J, Tohen M. Comparison of olanzapine and risperidone in the EMBLEM Study: translation of randomized controlled trial findings into clinical practice.. *Int Clin Psychopharmacol* 2010 ;25 (5) :257-63 Factor d'impacte: 3,345 Q2
- Novick D, Haro JM, Bertsch J, Haddad PM. Incidence of Extrapiramidal Symptoms and Tardive Dyskinesia in Schizophrenia: Thirty-Six-Month Results From the European Schizophrenia Outpatient Health Outcomes Study.. *J Clin Psychopharmacol* 2010 ;30 (5) :531-40 Factor d'impacte: 5,092 Q1
- Novick D, Haro JM, Suárez D, Perez V, Dittmann R, Haddad PM. Predictors and clinical consequences of non-adherence with antipsychotic medication in the outpatient treatment of schizophrenia.. *Psychiatry Res* 2010 ;176 (2-3) :109-13 Factor d'impacte: 2,373 Q2
- Ramírez N, Arranz B, Salavert J, Alvarez E, Corripio I, Dueñas RM, Perez V, San L. Predictors of schizophrenia in patients with a first episode of psychosis. *Psychiatry Res* 2010 ;175 (1-2) :11-4 Factor d'impacte: 2,666 Q2
- Reed C, Goetz I, Vieta E, Bassi M, Haro JM. Work impairment in bipolar disorder patients - results from a two-year observational study (EMBLEM).. *Eur J Psychiatry* 2010;25 (6) :338-44 Factor d'impacte: 0,462 Q4
- Rosa AR, Cruz N, Franco C, Haro JM, Bertsch J, Reed C, Aarre TF, Sánchez Moreno J, Vieta E. Why do clinicians maintain antidepressants in some patients with acute mania? hints from the european Mania in Bipolar Longitudinal Evaluation of Medication (EMBLEM), a large naturalistic study.. *J Clin Psychiatry* 2010 ;71 (8) :1000-6 Factor d'impacte: 5,218 Q1
- San L, Arranz B. Agomelatina: un nuevo enfoque antidepresivo para una remisión de calidad.. *Rev Psiquiatr Salud Mental* 2010 ;3 (Supl.1) :15-20 Q0
- Scott KM, Hwang I, Chiu WT, Kessler RC, Sampson NA, Angermeyer MC, Beautrais A, Borges G, Bruffaerts R, De Graaf R, Florescu S, Fukao A, Haro JM, Hu C, Kovess V, Levinson D, Posada Villa J, Scocco P, Nock MK. Chronic Physical Conditions and Their Association With First Onset of Suicidal Behavior in the World Mental Health Surveys.. *Psychol Med* 2010 ;72 (7) :712-9 Factor d'impacte: 5,012 Q1
- Scott KM, Wells JE, Angermeyer MC, Brugha TS, Bromet EJ, Demyttenaere K, De Girolamo G, Gureje O, Haro JM, Jin R, Karam A, Kovess V, Lara C, Levinson D, Ormel J, Posada Villa J, Sampson N, Takeshima T, Zhang M, Kessler RC. Gender and the relationship between marital status and first onset of mood, anxiety and substance use disorders.. *Psychol Med* 2010 ;40 (9) :1495-505 Factor d'impacte: 5,012 Q1
- Sevilla Dedieu C, Kovess V, Haro JM, Fernández A, Vilagut G, Alonso J. ESEMed/MHEDEA 2000 investigators. Seeking Help for Mental Health Problems Outside the Conventional Health Care System: Results From the European Study of the Epidemiology of Mental Disorders (ESEMed).. *Can J Psychiatry* 2010 ;55 (9) :586-97 Factor d'impacte: 2,293 Q2
- Stegmann ME, Ormel J, De Graaf R, Haro JM, De Girolamo G, Demyttenaere K, Kovess V, Matschinger H, Vilagut G, Alonso J, Burger H. ESEMed/MHEDEA 2000 investigators. Functional disability as an explanation of the associations between chronic physical conditions and 12-month major depressive episode.. *J Affect Disord* 2010 ;124 (1-2) :38-44 Factor d'impacte: 3,763 Q1
- Stein DJ, Scott K, Haro JM, Aguilar S, Alonso J, Angermeyer MC, Demyttenaere K, De Girolamo G, Iwata N, Posada Villa J, Kovess V, Lara C, Ormel J, Kessler RC, von Korff M. Early

childhood adversity and later hypertension: data from the World Mental Health Survey.. *Ann Clin Psychiatry* 2010 ;22 (1) :19-28 Q0